

# Paired Binary Data

ST551 Lecture 23

---

Charlotte Wickham

2017-11-15

**Finish last time's slides**

---

## Paired Binary Data

---

## Paired Binary Data

Imagine now that our two samples of Bernoulli populations aren't independent, but paired in some way.

$$Y_1, \dots, Y_n \sim \text{Bernoulli}(p_Y)$$

$$X_1, \dots, X_n \sim \text{Bernoulli}(p_X)$$

but  $(Y_i, X_i)$  are paired.

Examples:

- $n$  subjects with a disease, and  $n$  without a disease are sampled then matched (based on demographic factors), response is presence of some risk factor
- Sibling (or twin) studies:  $n$  pairs of related people where one falls in one group, and the other falls in the other group, observe some binary response on every person.
- Binary before and after measurements on the same person

## Paired Binary Data

Gather sample of  $n = 40$  voters.

Before debate: Will you vote for candidate A?

After debate: Will you vote for candidate A?

subject	before	after
1	1	1
2	1	0
3	1	0
4	1	1
5	1	1
6	1	0

## Just a $2 \times 2$ table?

---

	0	1
<b>after</b>	21	19
<b>before</b>	23	17

---

---

	0	1
<b>0</b>	12	11
<b>1</b>	9	8

---

## How to analyse?

**Option 1:** Treat like paired two sample data and do a paired t-test

**Option 2:** McNemar's test

## Paired t-test

Null hypothesis:  $H_0 : p_{\text{before}} = p_{\text{after}}$

Look at (per voter) differences:

subject	before	after	diff
1	1	1	0
2	1	0	-1
3	1	0	-1
4	1	1	0
5	1	1	0
6	1	0	-1

-1	0	1
9	20	11



## Paired t-test calculations

$$\bar{D} = \frac{1}{n} ((-1 \times 9) + (0 \times 20) + (1 \times 11)) = \frac{b - c}{n} = \frac{2}{40} = 0.05$$

$$\begin{aligned} s_D^2 &= \frac{1}{n-1} (9(-1 - \bar{D})^2 + 20(0 - \bar{D})^2 + 11(1 - \bar{D})^2) \\ &= \frac{1}{n-1} \left( c + b - \frac{(b-c)^2}{n} \right) \\ &= \frac{1}{40-1} \left( 9 + 11 - \frac{(11-9)^2}{n} \right) \\ &= 0.51 \end{aligned}$$

## Paired t-test calculations

```
##  
## One Sample t-test  
##  
## data:  df$diff  
## t = 0.4427, df = 39, p-value = 0.6604  
## alternative hypothesis: true mean is not equal to 0  
## 95 percent confidence interval:  
## -0.1784514  0.2784514  
## sample estimates:  
## mean of x  
##      0.05
```

## McNemar's test

Null hypothesis:  $H_0 : p_{\text{before}} = p_{\text{after}}$

Conditions on the number of discordant pairs,  $b + c$ .

	0	1
0	12	11
1	9	8

Under Null hypothesis, we expect the number of discordant pairs (e.g. people who change their minds during debate) should be equally split between  $b$  and  $c$ .

## McNemar's test

Conditional on  $b + c$ ,

$$b \sim \text{Binomial}(b + c, 0.5)$$

Do, one sample Z-test for proportions, leads to

$$Z = \frac{b - c}{\sqrt{b + c}} \sim N(0, 1) \quad \text{under null hypothesis}$$

(sometimes people square this statistic, and compare to  $\chi_1^2$ )

## Example: McNemar's

	0	1
0	12	11
1	9	8

$$Z = \frac{b - c}{\sqrt{b + c}} = \frac{11 - 9}{\sqrt{11 + 9}} = 0.45$$

Compare to  $N(0,1)$

- McNemar's test is equivalent to the paired t-test, in the sense that the two test statistics are monotone transformations of each other.
- For large sample sizes, the two test statistics get closer and closer to the same value: asymptotically equivalent.